

Auburn Montgomery Warhawk Health Services

7461 East Dr STE 102
Montgomery AL, 36117-8092
(334) 244-3281

Date: 03/26/24

PATIENT: OLIPHINT, DONNA **DOB:** 04/28/1951 **Gender:** FEMALE **MRN:** 1032 **AGE:** 72 Years

VISIT TYPE: Office Visit Established

CHIEF COMPLAINT: Travel consult to Luanda, Kenya

ALLERGIES: Diflucan ((Rash) or (C/O a rash))

MEDICATIONS:

Myrbetriq 25 mg tablet,extended release
acyclovir 400 mg tablet
atorvastatin 20 mg tablet
azelastine 137 mcg (0.1 %) nasal spray aerosol
bimatoprost 0.03 % eye drops
diltiazem CD 120 mg capsule,extended release 24 hr (TAKE ONE CAPSULE BY MOUTH ONE TIME DAILY)
estradiol 0.01% (0.1 mg/gram) vaginal cream
estradiol 2 mg tablet
fluticasone propionate 50 mcg/actuation nasal spray,suspension
hydrocortisone 2.5 % topical ointment
levothyroxine 112 mcg tablet

PAST MEDICAL HISTORY: hypothyroidism, HLD, overactive bladder, cardiac arrhythmia

PAST SURGICAL HISTORY: Hysterectomy

ORDERS:

TDap (1 unit(s)): Immunizations
Immunization Injection Administration (1 unit(s)): Injection

RESULTS:

TDap:
Site Given: Left Deltoid
Dosage Given: 0.5 milliliter(s)

ASSESSMENT:

Z23 - Encounter for immunization (PRIMARY).

PLAN:

Rx for malaria prevention. Rx in case she has traveler's diarrhea. Other recommended vaccines- yellow fever and check on polio vaccine as a child. She did not bring a copy of her yellow card but will let us know if these are vaccines she is in need of.

Cipro 500 mg tablet
Refills: 0 Qty: 14 TABLET(S)
Instructions: TAKE 1 TABLET (500 MG) BY ORAL ROUTE EVERY 12 HOURS for traveler's diarrhea.

ondansetron HCl 4 mg tablet
Refills: 0 Qty: 9 TABLET(S)
Instructions: TAKE 1 tablet PO every 6-8 hrs as needed for nausea/vomiting

doxycycline hyclate 100 mg tablet

Auburn Montgomery Warhawk Health Services

7461 East Dr STE 102
Montgomery AL, 36117-8092
(334) 244-3281

Date: 03/26/24

PATIENT: OLIPHINT, DONNA **DOB:** 04/28/1951 **Gender:** FEMALE **MRN:** 1032 **AGE:** 72 Years

Refills: 0 Qty: 45 TABLET(S)

Instructions: TAKE 1 TABLET (100 MG) BY ORAL ROUTE once daily for malaria prevention. Start 2 days prior to exposure and continue through 4 weeks after last day of exposure.

Electronically signed by Lindsey Barton, CRNP on 03/26/24 4:12 PM

L Barton, CRNP