

Carol Stewart, C.R.N.P.
Lic. #1-045437
Rx: 1178

AUBURN MONTGOMERY
STUDENT HEALTH CENTER
P. O. Box 244023
Montgomery, AL 36124-4023
(334) 244-3281

Ramona Lazenby, C.R.N.P.
Lic. #1-039092
Rx: 0703

Darlene Pierce, C.R.N.P.
Lic. #1-134315
RX: 3692

Maryluz Fuentes, M.D.
(334) 286-3444
DEA #BF 6451520
AL #24127

Name Charles Bush Date 7/20/17


Address _____

Rx

Malarone 250/100mg

Disp: 24

Sig: Take one pill a day. Begin 2 days before
departure, continue through stay and
for 7 days after return.

 M.D. or C.R.N.P.

Product Selection Permitted

Label all prescriptions

Refill: 1 2 3 5 PRN NR

Dispense As Written

M.D. or C.R.N.P.

medications?

Some other drugs can interact with atovaquone-proguanil and cause you problems. Your doctor is responsible for evaluating the other medicines you are taking to ensure that there are no interactions between them and atovaquone-proguanil. In some instances, medicines can be adjusted to minimize the interaction. You can also ask your pharmacist to check for drug interactions.

For example, people taking the common blood-thinning medicine called Coumadin may be at an increased risk for bleeding if they take atovaquone-proguanil. The Coumadin dose can be adjusted allowing you to take both medicines safely.

What are the potential side effects of atovaquone-proguanil?

Atovaquone-proguanil is well tolerated, and side effects are rare. The most common adverse reactions reported are stomach pain, nausea, vomiting, and headache. These side effects can often be lessened by taking atovaquone-proguanil with food.

All medicines may have some side effects. Minor side effects such as nausea, occasional vomiting, or diarrhea usually do not require stopping the antimalarial drug. If you cannot tolerate your antimalarial drug, see your health care provider; other antimalarial drugs are available.

neTM)

tions

travelers because the
ys before traveling to
a transmission occurs

ter trips because you
medicine for 7 days
than 4 weeks

sensitive than some of
pecially for trips of

ife to use oguanil?

CDC has no limits on the use of atovaquone-proguanil for the prevention of malaria. There is no evidence of harm when the drug has been used for extended periods of time.

For more information:

Check out the CDC malaria website at www.cdc.gov/malaria

Health-care providers needing assistance with diagnosis or management of suspected cases of malaria should call the CDC Malaria Hotline: 770-488-7788 or 855-856-4713 toll-free (M-F, 9am-5pm, eastern time).

Emergency consultation after hours, call: 770-488-7100 and request to speak with a CDC Malaria Branch clinician.

Prevent Malaria

- Take an antimalarial drug.
- Prevent mosquito bites.
- If you get sick, immediately seek professional medical care.