

Carol Stewart, C.R.N.P.
Lic. #1-044437
Rx: 1178

AUBURN MONTGOMERY
STUDENT HEALTH CENTER
P. O. Box 244023
Montgomery, AL 36124-023
(334) 244-3281

Ramona Lazenby, C.R.N.P.
Lic. #1-039092
Rx: 0703

Darlene Pierce, C.R.N.P.
Lic. #1-134315
RX: 3692

Maryluz Fuentes, M.D.
(334) 286-3444
DEA #BF 6451520
AL #24127

Name Mary Adams Date 6/2/16

Address _____

R

Vivotif
Disp: 4
Sig: Take as directed.

[Signature] M.D. or C.R.N.P.

Product Selection Permitted

Label all prescriptions

Refill: 1 2 3 5 PRN NR

Dispense As Written

Time to take your
2nd vaccine pill.
Take with cold or
lukewarm water
& 1 hour before a
meal. Put remaining
doses in fridge.

Free Text Message Reminders*

- Send a text message with the word "TRAVEL" to 21756 on the day you start Vivotif
- You'll receive text message reminders when it's time to take your remaining capsules
- This recurring program includes a total of 4 messages over the course of 7 days

***Notice:** The information requested will be used by PaxVax, Inc. to send text message reminders. Message and data rates may apply. At any time during the program, you may text "HELP" for assistance, or "STOP" to discontinue receiving messages. Full Terms and Conditions can be accessed at <http://mpulsemobile.com/paxvax-terms-conditions-21756/>.

Indications and Usage

Vivotif® (Typhoid Vaccine Live Oral Ty21a) is indicated for immunization of adults and children greater than 6 years of age against disease caused by *Salmonella* Typhi. Routine typhoid vaccination is not recommended in the United States of America. Selective immunization against typhoid fever is recommended for the following groups: 1) travelers to areas in which there is a recognized risk of exposure to *S. Typhi*, 2) persons with intimate exposure (e.g. household contact) to a *S. Typhi* carrier, and 3) microbiology laboratory workers who work frequently with *S. Typhi*. There is no evidence to support the use of typhoid vaccine to control common source outbreaks, disease following natural disasters or in persons attending rural summer camps.

Please see reverse side for additional
Important Safety Information.

 **Vivotif®**
Typhoid Vaccine Live Oral Ty21a

OVERDOSE: If overdose is suspected,

US residents can call the US National Poison Hotline at 1-800-222-1222. Canada residents can call a provincial poison control center.

ORAL (TIE-foyd)

y a certain bacteria
inking contaminated
ing to areas where
Asia), who are exposed
boratory. Typhoid
body to produce its own

th care provider before
Before taking each
pen the blister pack and
cist if the foil blister pack or
y every other day for 4 doses or
warm liquid about 1 hour before
mediately. Do not chew, crush, or
ne vaccine to be most effective.
your next dose. You should
posure to typhoid fever. You may
years. Ask your doctor or

all your doctor or pharmacist
because he or she has judged that
ing this medication do not have
however, seek immediate medical
ding: rash, itching/swelling
g. This is not a complete list of
your doctor or pharmacist.
numbers do not provide medical
se Event Reporting System
t 1-866-234-2345.

or emergency room immediately.