

Patient Information	Specimen Information	Client Information
FAUST, WANDA DOB: 07/02/1952 AGE: 66 Gender: F Fasting: Y Phone: NG Patient ID: NG	Specimen: AL525571A Requisition: 1919035 Collected: 01/15/2019 / 08:20 CST Received: 01/16/2019 / 03:02 CST Reported: 01/16/2019 / 05:45 CST	Client #: 49100638 QATL0000 TUCK, PAMELA P AUM NURSING CARE CENTER 7461 EAST DRIVE #102 MONTGOMERY, AL 36117

Test Name	In Range	Out Of Range	Reference Range	Lab
LIPID PANEL, STANDARD				
CHOLESTEROL, TOTAL	158		<200 mg/dL	AT
HDL CHOLESTEROL	52		>50 mg/dL	AT
TRIGLYCERIDES	124		<150 mg/dL	AT
LDL-CHOLESTEROL	83		mg/dL (calc)	AT

Reference range: <100

Desirable range <100 mg/dL for primary prevention;
<70 mg/dL for patients with CHD or diabetic patients
with > or = 2 CHD risk factors.

LDL-C is now calculated using the Martin-Hopkins
calculation, which is a validated novel method providing
better accuracy than the Friedewald equation in the
estimation of LDL-C.

Martin SS et al. JAMA. 2013;310(19): 2061-2068
(<http://education.QuestDiagnostics.com/faq/FAQ164>)

CHOL/HDL-C RATIO	3.0	<5.0 (calc)	AT
NON HDL CHOLESTEROL	106	<130 mg/dL (calc)	AT

For patients with diabetes plus 1 major ASCVD risk
factor, treating to a non-HDL-C goal of <100 mg/dL
(LDL-C of <70 mg/dL) is considered a therapeutic
option.

COMPREHENSIVE METABOLIC PANEL				AT
GLUCOSE	97		65-99 mg/dL	

Fasting reference interval

UREA NITROGEN (BUN)	15	7-25 mg/dL
CREATININE	0.70	0.50-0.99 mg/dL

For patients >49 years of age, the reference limit
for Creatinine is approximately 13% higher for people
identified as African-American.

eGFR NON-AFR. AMERICAN	90	> OR = 60 mL/min/1.73m ²	
eGFR AFRICAN AMERICAN	105	> OR = 60 mL/min/1.73m ²	
BUN/CREATININE RATIO	NOT APPLICABLE	6-22 (calc)	
SODIUM	138	135-146 mmol/L	
POTASSIUM	4.2	3.5-5.3 mmol/L	
CHLORIDE	103	98-110 mmol/L	
CARBON DIOXIDE	22	20-32 mmol/L	
CALCIUM	9.9	8.6-10.4 mg/dL	
PROTEIN, TOTAL	7.3	6.1-8.1 g/dL	
ALBUMIN	4.4	3.6-5.1 g/dL	
GLOBULIN	2.9	1.9-3.7 g/dL (calc)	
ALBUMIN/GLOBULIN RATIO	1.5	1.0-2.5 (calc)	
BILIRUBIN, TOTAL	0.7	0.2-1.2 mg/dL	
ALKALINE PHOSPHATASE	89	33-130 U/L	
AST	14	10-35 U/L	
ALT	18	6-29 U/L	
HEMOGLOBIN A1c	5.6	<5.7 % of total Hgb	AT

For the purpose of screening for the presence of

Patient Information	Specimen Information	Client Information
FAUST, WANDA DOB: 07/02/1952 AGE: 66 Gender: F Fasting: Y Patient ID: NG	Specimen: AL525571A Collected: 01/15/2019 / 08:20 CST Received: 01/16/2019 / 03:02 CST Reported: 01/16/2019 / 05:45 CST	Client #: 49100638 TUCK, PAMELA P

Test Name	In Range	Out Of Range	Reference Range	Lab
diabetes:				

<5.7% 5.7-6.4% > or =6.5%	Consistent with the absence of diabetes Consistent with increased risk for diabetes (prediabetes) Consistent with diabetes
---------------------------------	---

This assay result is consistent with a decreased risk of diabetes.

Currently, no consensus exists regarding use of hemoglobin A1c for diagnosis of diabetes in children.

According to American Diabetes Association (ADA) guidelines, hemoglobin A1c <7.0% represents optimal control in non-pregnant diabetic patients. Different metrics may apply to specific patient populations. Standards of Medical Care in Diabetes (ADA).

PERFORMING SITE:

AT QUEST DIAGNOSTICS-ATLANTA, 1777 MONTREAL CIRCLE, TUCKER, GA 30084-6802 Laboratory Director: ANDREW N YOUNG, MD, PHD, CLIA: 11D0255931