

Carol Stewart, C.R.N.P.
Lic. #1-045437
Rx: 1178

AUBURN MONTGOMERY
STUDENT HEALTH CENTER
P.O. Box 244023
Montgomery, AL 36124-4023
(334) 244-3281

Ramona Lazenby, C.R.N.P.
Lic. #1-039092
Rx: 0703

Darlene Pierce, C.R.N.P.
Lic. #1-134315
RX: 3692


Maryluz Fuentes, M.D.
(334) 286-3444
DEA #BF 6451520
AL #24127

COT-V17-CVRS RV (12-09)
RP (4-1-11)

Room Mirror

ER

Vaccine



used

own

Name Makenzie Anthony Date 3/29/16

Address _____

R

Vivotif

Disp: 4

Sig: Take as directed.

[Signature]

M.D. or C.R.N.P.

Label all prescriptions

Refill: 1 2 3 5 PRN (NR)

Product Selection Permitted

Dispense As Written

M.D. or C.R.N.P.

Time to take your
2nd vaccine pill.
Take with cold or
lukewarm water
& 1 hour before a
meal. Put remaining
doses in fridge.

Free Text Message Reminders*

- Send a text message with the word "TRAVEL" to 21756 on the day you start Vivotif
- You'll receive text message reminders when it's time to take your remaining capsules
- This recurring program includes a total of 4 messages over the course of 7 days

*Notice: The information requested will be used by PaxVax, Inc. to send text message reminders. Message and data rates may apply. At any time during the program, you may text "HELP" for assistance, or "STOP" to discontinue receiving messages. Full Terms and Conditions can be accessed at <http://mpulsemobile.com/paxvax-terms-conditions-21756/>.

Indications and Usage

Vivotif® (Typhoid Vaccine Live Oral Ty21a) is indicated for immunization of adults and children greater than 6 years of age against disease caused by *Salmonella* Typhi. Routine typhoid vaccination is not recommended in the United States of America. Selective immunization against typhoid fever is recommended for the following groups: 1) travelers to areas in which there is a recognized risk of exposure to *S. Typhi*, 2) persons with intimate exposure (e.g. household contact) to a *S. Typhi* carrier, and 3) microbiology laboratorians who work frequently with *S. Typhi*. There is no evidence to support the use of typhoid vaccine to control common source outbreaks, disease following natural disasters or in persons attending rural summer camps.

Please see reverse side for additional
Important Safety Information.



Vivotif®

Typhoid Vaccine Live Oral Ty21a

health care provider before
r. Before taking each
y open the blister pack and
pharmacist if the foil blister pack or
lly every other day for 4 doses or
ly warm liquid about 1 hour before
mediately. Do not chew, crush, or
the vaccine to be most effective.
ke your next dose. You should
posure to typhoid fever. You may
5 years. Ask your doctor or

tell your doctor or pharmacist
because he or she has judged that
using this medication do not have
. However, seek immediate medical
luding: rash, itching/swelling
ing. This is not a complete list of
act your doctor or pharmacist.
ng numbers do not provide medical
arse Event Reporting System
a at 1-866-234-2345.

OVERDOSE: If overdose is suspected, contact a poison control center or emergency room immediately. US residents can call the US National Poison Hotline at 1-800-222-1222. Canada residents can call a provincial poison control center.