

Carol Stewart, C.R.N.P.
Lic. #1-045437
Rx: 1178

ADRIAN MONTGOMERY
STUDENT HEALTH CENTER
P. O. Box 244023
Montgomery, AL 36124-4023
(334) 244-3281

Ramon Zenby, C.R.N.P.
Lic. #039092
Rx: 0703

Darlene Pierce, C.R.N.P.
Lic. #1-134315
RX: 3692

Maryluz Fuentes, M.D.
(334) 286-3444
DEA #BF 6451520
AL #24127


RAL (TIE-foyd)

Name Tammy Hudson Date 9/16/16

Address _____

Rx

Vivotif
Disp: 4
Sig: Take as directed.


Product Selection Permitted M.D. or C.R.N.P.

Label all prescriptions

Refill: 1 2 3 5 PRN NR

Dispense As Written M.D. or C.R.N.P.

where there is a risk of exposure (e.g., contact) to someone who carries *Salmonella* Typhi (STyphi), the bacterium that causes typhoid fever, and people who work in a laboratory with STyphi.

Not everyone who takes Vivotif will be fully protected against typhoid fever. Therefore, you should continue to take precautions against typhoid fever exposure. Vivotif does not protect against other *Salmonella* species or other bacteria that cause disease of the gastrointestinal tract. Vivotif should not be used to treat infections with STyphi.

Important Safety Information

Do not take Vivotif if:

- You are allergic (hypersensitive) to any of the ingredients contained in Vivotif
- You have a poor immune system for any reason
- You have an infection with a high temperature
- You have an acute gastrointestinal illness

Tell your health care professional if you are taking or have recently taken any other medicines. This is because Vivotif can affect the way some medicines work and some medicines can affect the way Vivotif works. In particular, tell your health care professional if you are taking antibiotics or medicines to prevent malaria. Postpone taking Vivotif if persistent diarrhea or vomiting is occurring. It is essential that all 4 doses of vaccine be taken at the prescribed alternate day interval to obtain a maximal protective immune response. In clinical trials, the most common side effects observed were abdominal pain, nausea, headache, fever, diarrhea, vomiting, and skin rash.

Report any serious adverse reactions to your health care professional. You may also report an adverse reaction directly to the Vaccine Adverse Event Reporting System at 1-800-822-7967 or <http://vaers.hhs.gov>.

For more information about Vivotif, please see accompanying full Prescribing Information.

© 2016 PaxVax. All Rights Reserved. | PaxVax, Inc., 500 Veterans Boulevard, Suite 500, Redwood City, CA 94063
PAX-US-0035 | FEB 2016

PaxVax

attention if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. Contact your doctor for medical advice about side effects. The following numbers do not provide medical advice, but in the US, you may report side effects to the Vaccine Adverse Event Reporting System (VAERS) at 1-800-822-7967. In Canada, you may call Health Canada at 1-866-234-2345.

OVERDOSE: If overdose is suspected, contact a poison control center or emergency room immediately. US residents can call the US National Poison Hotline at 1-800-222-1222. Canada residents can call a provincial poison control center.

a certain bacteria making contaminated g to areas where Asia), who are exposed oratory.Typhoid ody to produce its own

th care provider before Before taking each open the blister pack and e foil blister pack or

bt damaged. Consult your ion by mouth without food, usually every other day for 4 doses or capsule whole with cold or barely warm liquid about 1 hour before and swallow with the liquid immediately. Do not chew, crush, or the dosing schedule closely for the vaccine to be most effective. idar to keep track of when to take your next dose. You should east 1 week before possible exposure to typhoid fever. You may n at risk for typhoid fever after 5 years. Ask your doctor or

this effect persists or worsens, tell your doctor or pharmacist as prescribed this medication because he or she has judged that of side effects. Many people using this medication do not have gic reaction to this drug is rare. However, seek immediate medical

Carol Stewart, C.R.N.P.
Lic. #1-045437
Rx: 1178

AUBURN MONTGOMERY
STUDENT HEALTH CENTER
P. O. Box 244023
Montgomery, AL 36124-4023
(334) 244-3281

Ramona Lazenby, C.R.N.P.
Lic. #1-039092
Rx: 0703

Darlene Pierce, C.R.N.P.
Lic. #1-134315
RX: 3692

Maryluz Fuentes, M.D.
(334) 286-3444
DEA #BF 6451520
AL #24127

Name Daniel Hudson Date 9/16/16
Address _____

Rx Malarone 250/100mg

Disp: 19

Sig: Take one pill a day. Begin 2 days before departure, continue through stay and for 7 days after return.

[Signature]
Product Selection Permitted

Label all prescriptions

Refill: 1 2 3 5 PRN NR

Dispense As Written

M.D. or C.R.N.P.

en depends on their weight.
aily dose for children based

e a ½ pediatric tablet daily,
et daily
let daily
lets daily
lets daily

ould be taken with food or a

itovaquone-

Antimalarial drugs are available in the United States by prescription only. Medicines should be obtained at a pharmacy before travel rather than in the destination as abroad has its risks: quality, contaminated, or t you against malaria.

Carol Stewart, C.R.N.P.
Lic. #1-045437
Rx: 1178

AUBURN MONTGOMERY
STUDENT HEALTH CENTER
P. O. Box 244023
Montgomery, AL 36124-4023
(334) 244-3281

Ramona Lazenby, C.R.N.P.
Lic. #1-039092
Rx: 0703

Darlene Pierce, C.R.N.P.
Lic. #1-134315
RX: 3692

Maryluz Fuentes, M.D.
(334) 286-3444
DEA #BF 6451520
AL #24127

Name Tammy Hudson Date _____
Address _____

Rx Malarone 250/100mg

Disp: 19

Sig: Take one pill a day. Begin 2 days before departure, continue through stay and for 7 days after return.

[Signature]
Product Selection Permitted

Label all prescriptions

Refill: 1 2 3 5 PRN NR

Dispense As Written

M.D. or C.R.N.P.

he world can
uanil be used for
aria in travelers?

uanil can be taken in any
alaria occurs.

laces in the world where
s and the malaria drugs
use in each place. This
using the malaria map on
ww.cdc.gov/malaria/map/



This map shows an approximation of the parts of the world where malaria transmission occurs. For more detailed information about the occurrence of malaria transmission in specific countries, please use the interactive Malaria Map Application.

Before traveling to the area where malaria transmission occurs. They should take one dose per day while there, and for 7 consecutive days after leaving. In rare instances (such as if you switched to atovaquone-proguanil from another malaria medicine) your doctor may instruct you to take it for a longer period of time.

The daily dosage for adults is a single adult tablet.

Center for Global Health
Division of Parasitic Diseases and Malaria

