

Carol Stewart, C.R.N.P.
Lic. #1-045437
Rx: 1178

AUBURN MONTGOMERY
STUDENT HEALTH CENTER
P. O. Box 244023
Montgomery, AL 36124-4023
(334) 244-3281

Ramona Lazenby, C.R.N.P.
Lic. #1-039092
Rx: 0703

Darlene Pierce, C.R.N.P.
Lic. #1-134315
RX: 3692

Maryluz Fuentes, M.D.
(334) 286-3444
DEA #BF 6451520
AL #24127

Name Faye Ageni

Address _____

Date 29 Nov 2016

R

Malarone 250/100mg

Disp: 23

Sig: Take one pill a day. Begin 2 days before departure, continue through stay and for 7 days after return.

[Signature], M.D. or C.R.N.P.
Product Selection Permitted

Label all prescriptions

Refill: 1 2 3 5 PRN NR

Dispense As Written

M.D. or C.R.N.P.



Who can take atovaquone-proguanil?

Atovaquone-proguanil can be prescribed to adults and children who weigh at least 11 pounds (5 kg).

Who should not take atovaquone-proguanil?

Pregnant women, women who are nursing infants less than 5kg, or infants who weigh less than 5kg should not take atovaquone-proguanil. Also, people with severe kidney disease should not take atovaquone-proguanil.

How should I take atovaquone-proguanil?

Both adults and children should take one dose of atovaquone-proguanil per day starting a day or two before traveling to the area where malaria transmission occurs. They should take one dose per day while there, and for 7 consecutive days after leaving. In rare instances (such as if you switched to atovaquone-proguanil from another malaria medicine) your doctor may instruct you to take it for a longer period of time.

The daily dosage for adults is a single adult tablet.

depends on their weight.
/ dose for children based

1/2 pediatric tablet daily,
daily
daily
daily
daily

should be taken with food or a

atovaquone-

available in the United States by
should be obtained at a

pharmacy before travel rather than in the destination country. Buying medications abroad has its risks: the drugs could be of poor quality, contaminated, or counterfeit and not protect you against malaria.

In what parts of the world can atovaquone-proguanil be used for prevention of malaria in travelers?

Currently atovaquone-proguanil can be taken in any part of the world where malaria occurs.

CDC keeps track of all the places in the world where malaria transmission occurs and the malaria drugs that are recommended for use in each place. This information can be found using the malaria map on the CDC website: <http://www.cdc.gov/malaria/map/index.html>.

