

Carol Stewart, C.R.N.P.
Lic. #1-04543
Rx: 1178

AUBURN MONTGOMERY
STUDENT HEALTH CENTER
P. O. Box 244023
Montgomery, AL 36124-4023
(334) 244-3281

Ramona Lazenby, C.R.N.P.
Lic. #1-039092
Rx: 0703

Darlene Pierce, C.R.N.P.
Lic. #1-134315
RX: 3692

Maryluz Fuentes, M.D.
(334) 286-3444
DEA #BF 6451520
AL #24127

Traveling

Name Susan Hunt

Date 2/20/18

Address _____

ren depends on their weight.
daily dose for children based

R

Malarone 250/100mg

Disp: 16

Sig: Take one pill a day. Begin 2 days before
departure, continue through stay and
for 7 days after return.

ke a ½ pediatric tablet daily,
plet daily
blet daily
blets daily
blets daily
ily
ould be taken with food or a

L. Fuentes, M.D. or C.R.N.P.

Label all prescriptions


Product Selection Permitted

Refill: 1 2 3 5 PRN NR

Dispense As Written

M.D. or C.R.N.P.

atovaquone-



Antimalarial drugs are available in the United States by
prescription only. Medicines should be obtained at a
pharmacy before travel rather than in the destination
country. Buying medications abroad has its risks:
the drugs could be of poor quality, contaminated, or
counterfeit and not protect you against malaria.

Who can take atovaquone-proguanil?

Atovaquone-proguanil can be prescribed to adults and
children who weigh at least 11 pounds (5 kg).

Who should not take atovaquone-proguanil?

Pregnant women, women who are nursing infants less
than 5kg, or infants who weigh less than 5kg should not
take atovaquone-proguanil. Also, people with severe
kidney disease should not take atovaquone-proguanil.

How should I take atovaquone-proguanil?


Both adults and children should take one dose of
atovaquone-proguanil per day starting a day or two
before traveling to the area where malaria transmission
occurs. They should take one dose per day while there,
and for 7 consecutive days after leaving. In rare instances
(such as if you switched to atovaquone-proguanil from
another malaria medicine) your doctor may instruct you
to take it for a longer period of time.

The daily dosage for adults is a single adult tablet.

In what parts of the world can atovaquone-proguanil be used for prevention of malaria in travelers?

Currently atovaquone-proguanil can be taken in any
part of the world where malaria occurs.

CDC keeps track of all the places in the world where
malaria transmission occurs and the malaria drugs
that are recommended for use in each place. This
information can be found using the malaria map on
the CDC website: <http://www.cdc.gov/malaria/map/index.html>.



Center for Global Health
Division of Parasitic Diseases and Malaria

