

Carol Stewart, C.R.N.P.
Lic. #1045437
Rx: 1178

AUBURN MONTGOMERY
STUDENT HEALTH CENTER
P. O. Box 244023
Montgomery, AL 36124-4023
(334) 244-3281

Ramona Lazenby, C.R.N.P.
Lic. #1-039092
Rx: 0703

Darlene Pierce, C.R.N.P.
Lic. #1-134315
RX: 3692

Maryluz Fuentes, M.D.
(334) 286-3444
DEA #BF 6451520
AL #24127

Name Sarah Bragg Date 3/12/15

Address:

Rx

Doxycycline 100mg

Disp: 39

Sig: One pill a day; begin 2 days before
departure, continue through stay and
for four weeks after return.

Label all prescriptions

Refill: 1 2 3 5 PRN NR


Product Selection Permitted

M.D. or C.R.N.P.

Dispense As Written

M.D. or C.R.N.P.

for the prevention of malaria infection associated
with travel.



Who can take doxycycline?

Doxycycline can be prescribed to adults and children who are at least 8 years old.

Who should not take doxycycline?

It should not be used by pregnant women. It should not be given to children under the age of 8 years.

How should I take doxycycline?

Both adults and children should take one dose of doxycycline per day starting a day or two before traveling to the area where malaria transmission occurs. They should take one dose per day while there, and for 28 consecutive days after leaving.

The daily dosage for adults is 100mg per day.

Your doctor will have calculated the correct daily dose for your child based on the child's weight. The child's dose should not exceed the adult dose of 100mg per day.

Traveling

ne on a full stomach with a full
milk and other dairy products
urs of taking your doxycycline
vn for 1 hour after taking
nt irritation in your throat.

Why doxycycline?

re available in the United States
Medicines should be obtained
e travel rather than in the
Buying medications abroad
gs could be of poor quality,
interfeit and not protect you

Will doxycycline interact with my other medications?

Some other drugs can interact with doxycycline and cause you problems. Your doctor is responsible for evaluating the other medicines you are taking to ensure that there are no interactions between them and doxycycline. In some instances, medicines can be adjusted to minimize the interaction. You can also ask your pharmacist to check for drug interactions.

For example, people taking the common blood-thinning medicine called Coumadin (warfarin) may be at an increased risk for bleeding if they take doxycycline. If there are no alternatives, the Coumadin dose can be adjusted allowing you to take both medicines safely. Certain antacids (containing calcium or magnesium) should also be avoided while taking doxycycline.

Malaria is a serious disease that can cause death if not treated right away. It is caused by a parasite that can infect a certain type of mosquito which feeds on humans.

About 1,500 cases of malaria are diagnosed in the United States each year, almost all in travelers arriving from parts of the world where malaria occurs.

Center for Global Health
Division of Parasitic Diseases and Malaria

